

Ricardo Parreira da Silva , Universidade Estadual Paulista Júlio de Mesquita Filho, Brazil. email: MISSING

LOWER BOUNDS ON BLOW UP SOLUTIONS OF THE THREE-DIMENSIONAL NAVIER–STOKES EQUATIONS IN HOMOGENEOUS SOBOLEV SPACES

Suppose that $u(t)$ is a (weak) solution of the three-dimensional Navier–Stokes equations, either on the whole space or with periodic boundary conditions, that has a singularity at time T . We show that the norm of $u(T-t)$ in the homogeneous Sobolev space \dot{H}^s must be bounded below by $c_s t^{-(2s-1)/4}$ for $1/2 < s < 5/2$ ($s \neq 3/2$), where c_s is an absolute constant depending only on s ; and by $c_s \|u_0\|_{L^2}^{(5-2s)/5} t^{-2s/5}$ for $s > 5/2$. (The result for $1/2 < s < 3/2$ follows from well known lower bounds on blowup in L^p spaces.) We show in particular that the local existence time in $\dot{H}^s(\mathbb{R}^3)$ depends only on the \dot{H}^s -norm for $1/2 < s < 5/2$, $s \neq 3/2$.

Work in collaboration with: James Cooper Robinson (University of Warwick) and Witold Sadowski (Warsaw University).

Mathematical Reviews subject classification: Primary: ; Secondary:
Key words: ,